

Available online at www.sciencedirect.com**ScienceDirect**

AKCE International Journal of Graphs and Combinatorics 12 (2015) 64–69

www.elsevier.com/locate/akcej
AKCE
International
Journal of
Graphs and
Combinatorics

Harary index and some Hamiltonian properties of graphs

Rao Li

Department of mathematical sciences, University of South Carolina Aiken, Aiken, SC 29801, United States

Received 15 November 2013; accepted 19 May 2015

Available online 23 July 2015

Abstract

For a nontrivial connected graph G , its Harary index $H(G)$ is defined as

$$\sum_{\{u, v\} \subseteq V(G)} \frac{1}{d_G(u, v)},$$

where $d_G(u, v)$ is the distance between vertices u and v . Hua and Wang (2013), using Harary index, obtained a sufficient condition for the traceable graphs. In this note, we use Harary index to present sufficient conditions for Hamiltonian and Hamilton-connected graphs.

© 2015 Kalasalingam University. Production and Hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Keywords: Harary index; Hamiltonicity; Hamilton-connectivity

1. Introduction

We consider only finite undirected graphs without loops or multiple edges. Notation and terminology not defined here follow that in [1]. For a graph $G = (V, E)$, we use n and e to denote its order $|V|$ and size $|E|$, respectively. For two vertices u and v in a graph G , we use $d_G(u, v)$ to denote the distance between them. A cycle C in a graph G is called a Hamiltonian cycle of G if C contains all the vertices of G . A graph G is called Hamiltonian if G has a Hamiltonian cycle. A path P in a graph G is called a Hamiltonian path of G if P contains all the vertices of G . A graph G is called traceable if G has a Hamiltonian path. A graph G is called Hamilton-connected if for each pair of vertices in G there is a Hamiltonian path between them. If G and H are two vertex-disjoint graphs, we use $G \vee H$ to denote the join of G and H . We use $C(n, r)$ to denote the number of r -combinations of a set with n elements.

In 1993, Ivanciuc et al. [2] and Plavšić et al. [3] independently introduced the Harary index as a molecular descriptor. For a nontrivial connected graph G , its Harary index, denoted $H(G)$, is defined as

$$\sum_{\{u, v\} \subseteq V(G)} \frac{1}{d_G(u, v)}.$$

Peer review under responsibility of Kalasalingam University.

E-mail address: raol@usca.edu.

<http://dx.doi.org/10.1016/j.akcej.2015.06.010>

0972-8600/© 2015 Kalasalingam University. Production and Hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

In [4], Hua and Wang, using Harary index, obtained the following result for the traceable graphs.

Theorem 1 ([4]). *Let G be a connected graph of order $n \geq 4$. If*

$$H(G) \geq \frac{n^2 - 3n + 5}{2},$$

then G is traceable, unless $G = K_1 \vee (K_{n-3} \cup 2K_1)$, or $K_2 \vee (3K_1 \cup K_2)$, or $K_4 \vee 6K_1$.

In this note, using the ideas and techniques in [4], we will prove the following theorems on the Hamiltonian and Hamilton-connected graphs.

Theorem 2. *Let G be a connected graph of order $n \geq 3$. If*

$$H(G) \geq \frac{n^2 - 2n + 2}{2},$$

then G is Hamiltonian, unless $G = K_1 \vee (K_1 \cup K_{n-2})$ or $K_2 \vee (K_2^c \cup K_1)$.

Theorem 3. *Let G be a connected graph of order n . If*

$$H(G) \geq \frac{n^2 - 2n + 3}{2},$$

then G is Hamilton-connected, unless $G = K_2 \vee (K_1 \cup K_{n-3})$ or $K_3 \vee (3K_1)$.

Theorem 4. *Let $G = (X, Y; E)$, where $X = \{x_1, x_2, \dots, x_n\}$, $Y = \{y_1, y_2, \dots, y_n\}$, and $n \geq 2$, be a connected bipartite graph. If*

$$H(G) \geq \frac{9n^2 - 3n - 4}{6},$$

then G is Hamiltonian, unless $G = P_4$, a path having four vertices and three edges.

Theorem 5. *Let G be a 2-connected graph of order $n \geq 12$. If*

$$H(G) \geq \frac{n^2 - 3n + 7}{2},$$

then G is Hamiltonian or $G = K_2 \vee ((2K_1) \cup K_{n-4})$.

Theorem 6. *Let G be a 3-connected graph of order $n \geq 18$. If*

$$H(G) \geq \frac{n^2 - 4n + 15}{2},$$

then G is Hamiltonian or $G = K_3 \vee ((3K_1) \cup K_{n-6})$.

Theorem 7. *Let G be a k -connected graph of order n . If*

$$H(G) \geq \frac{2n(n-1) - (k+1)(n-k-1) + 1}{4},$$

then G is Hamiltonian.

2. Lemmas

In order to prove the theorems above, we need the following results as our lemmas.

Lemma 1. *Let G be a graph of order $n \geq 3$ with degree sequence $d_1 \leq d_2 \leq \dots \leq d_n$. If*

$$d_k \leq k < \frac{n}{2} \implies d_{n-k} \geq n - k,$$

then G is Hamiltonian.

Lemma 2. Let G be a graph of order $n \geq 3$ with degree sequence $d_1 \leq d_2 \leq \dots \leq d_n$. If

$$2 \leq k \leq \frac{n}{2}, \quad d_{k-1} \leq k \implies d_{n-k} \geq n - k + 1,$$

then G is Hamilton-connected.

Lemma 3. Let $G = (X, Y; E)$ be a bipartite graph such that $X = \{x_1, x_2, \dots, x_n\}$, $Y = \{y_1, y_2, \dots, y_n\}$, $n \geq 2$, and

$$\begin{aligned} d(x_1) &\leq d(x_2) \leq \dots \leq d(x_n), \\ d(y_1) &\leq d(y_2) \leq \dots \leq d(y_n). \end{aligned}$$

If

$$d(x_k) \leq k < n \implies d(y_{n-k}) \geq n - k + 1,$$

then G is Hamiltonian.

Lemma 4 ([5]). Let G be a 2-connected graph of order $n \geq 12$. If $e(G) \geq C(n-2, 2) + 4$, then G is Hamiltonian or $G = K_2 \vee ((2K_1) \cup K_{n-4})$.

Lemma 5 ([5]). Let G be a 3-connected graph of order $n \geq 18$. If $e(G) \geq C(n-3, 2) + 9$, then G is Hamiltonian or $G = K_3 \vee ((3K_1) \cup K_{n-6})$.

Lemma 6 ([5]). Let G be a k -connected graph of order n . If $e(G) \geq C(n, 2) - (k+1)(n-k-1)/2 + 1$, then G is Hamiltonian.

Notice that Lemma 1 is Corollary 3 on Page 209 in [6] or Theorem 4.5 on Page 57 in [1], Lemma 2 is Theorem 12 on Page 218 in [6], Lemma 3 is Corollary 5 on Page 210 in [6], and Lemmas 4–6 can be found in [5]. As in [4], we use $\widehat{D}_G(v)$ to denote

$$\sum_{u \in V(G)} \frac{1}{d_G(u, v)}.$$

Thus

$$\widehat{D}_G(v) \leq d(v) + \frac{1}{2}(n-1-d(v)).$$

Hence

$$\begin{aligned} H(G) &= \frac{1}{2} \sum_{v \in V(G)} \widehat{D}_G(v) \leq \frac{1}{2} \sum_{v \in V(G)} \left(d(v) + \frac{1}{2}(n-1-d(v)) \right) \\ &= \frac{1}{2} \sum_{v \in V(G)} \left(\frac{n-1}{2} + \frac{1}{2}d(v) \right) = \frac{n(n-1)}{4} + \frac{1}{4} \sum_{v \in V(G)} d(v). \end{aligned}$$

The above upper bound for $H(G)$ will be used in our proofs in Section 3.

3. Proofs

Proof of Theorem 2. Let G be a graph satisfying the conditions in Theorem 2. Suppose that G is not Hamiltonian. Then from Lemma 1, there exists an integer $k < \frac{n}{2}$ such that $d_k \leq k$ and $d_{n-k} \leq n - k - 1$. Obviously, $k \geq 1$. Therefore

$$\begin{aligned} H(G) &\leq \frac{n(n-1)}{4} + \frac{1}{4} \sum_{v \in V(G)} d(v) \\ &\leq \frac{n(n-1)}{4} + \frac{1}{4} \left(k^2 + (n-2k)(n-k-1) + k(n-1) \right) \end{aligned}$$

$$\begin{aligned}
&= \frac{n(n-1)}{4} + \frac{1}{2} + \frac{(n-1)(n-2)}{4} - \frac{(k-1)(2n-3k-4)}{4} \\
&= \frac{n^2-2n+2}{2} - \frac{(k-1)(k-2)}{4} - \frac{(k-1)(n-2k-1)}{2}.
\end{aligned}$$

From $H(G) \geq \frac{n^2-2n+2}{2}$, $k \geq 1$, and $n > 2k$, we have that $H(G) = \frac{n^2-2n+2}{2}$, $k = 1$ or $k = 2$ and $n = 2k + 1$, $d_1 = \dots = d_k = k$, $d_{k+1} = \dots = d_{n-k} = n - k - 1$, and $d_{n-k+1} = \dots = d_n = n - 1$.

If $k = 1$, then $d_1 = 1$, $d_2 = d_3 = \dots = d_{n-1} = n - 2$, and $d_n = n - 1$. Thus $G = K_1 \vee (K_1 \cup K_{n-2})$, which is not Hamiltonian.

If $k = 2$ and $n = 2k + 1$, we have $n = 5$. Therefore $d_1 = 2$, $d_2 = 2$, $d_3 = 2$, $d_4 = 4$, and $d_5 = 4$. Hence $G = K_2 \vee (K_2^c \cup K_1)$, which is not Hamiltonian.

This completes the proof of [Theorem 2](#). \square

Proof of Theorem 3. Let G be a graph satisfying the conditions in [Theorem 3](#). Suppose that G is not Hamilton-connected. Then from [Lemma 2](#), there exists an integer k with $2 \leq k \leq \frac{n}{2}$ such that $d_{k-1} \leq k$ and $d_{n-k} \leq n - k$. Therefore

$$\begin{aligned}
H(G) &\leq \frac{n(n-1)}{4} + \frac{1}{4} \sum_{v \in V(G)} d(v) \\
&\leq \frac{n(n-1)}{4} + \frac{1}{4} (k(k-1) + (n-2k+1)(n-k) + k(n-1)) \\
&= \frac{n(n-1)}{4} + 1 + \frac{(n-1)(n-2)}{4} - \frac{(k-2)(2n-3k-3)}{4} \\
&= \frac{n^2-2n+3}{2} - \frac{(k-2)(k-3)}{4} - \frac{(k-2)(n-2k)}{2}.
\end{aligned}$$

From $H(G) \geq \frac{n^2-2n+3}{2}$, $k \geq 2$, and $n \geq 2k$, we have that $H(G) = \frac{n^2-2n+3}{2}$, $k = 2$ or $(k = 3$ and $n = 2k)$, $d_1 = \dots = d_{k-1} = k$, $d_k = \dots = d_{n-k} = n - k$, and $d_{n-k+1} = \dots = d_n = n - 1$.

If $k = 2$, then $d_1 = 2$, $d_2 = d_3 = \dots = d_{n-2} = n - 2$, and $d_{n-1} = d_n = n - 1$. Thus $G = K_2 \vee (K_1 \cup K_{n-3})$, which is not Hamilton-connected.

If $k = 3$ and $n = 2k$, we have that $n = 6$. Therefore $d_1 = 3$, $d_2 = 3$, $d_3 = 3$, $d_4 = 5$, $d_5 = 5$, and $d_6 = 5$. Hence $G = K_3 \vee (3K_1)$, which is not Hamilton-connected.

This completes the proof of [Theorem 3](#). \square

Proof of Theorem 4. Let G be a graph satisfying the conditions in [Theorem 4](#). Suppose that G is not Hamiltonian. Then from [Lemma 3](#), there exists an integer $k < n$ such that $d(x_k) \leq k$ and $d(y_{n-k}) \leq n - k$. Next we will find an upper bound for $\widehat{D}_G(x_1)$. Let $N_G(x_1) := \{z_1, z_2, \dots, z_s\}$ be the neighbor of x_1 , where $s = d(x_1)$. Then $d_G(x_1, z_i) = 1$ for each $z_i \in N_G(x_1)$, $d_G(x_1, x_i) \geq 2$ for each x_i with $2 \leq i \leq n$, and $d_G(x_1, y_i) \geq 3$ for each $y_i \in Y - N_G(x_1)$. Thus

$$\widehat{D}_G(x_1) \leq d(x_1) + \frac{1}{2}(n-1) + \frac{1}{3}(n-d(x_1)) = \frac{5}{6}n - \frac{1}{2} + \frac{2}{3}d(x_1).$$

Similarly, we have that for each i with $2 \leq i \leq n$ and each j with $1 \leq j \leq n$,

$$\begin{aligned}
\widehat{D}_G(x_i) &\leq d(x_i) + \frac{1}{2}(n-1) + \frac{1}{3}(n-d(x_1)) = \frac{5}{6}n - \frac{1}{2} + \frac{2}{3}d(x_i), \\
\widehat{D}_G(y_j) &\leq d(y_j) + \frac{1}{2}(n-1) + \frac{1}{3}(n-d(y_j)) = \frac{5}{6}n - \frac{1}{2} + \frac{2}{3}d(y_j).
\end{aligned}$$

Therefore

$$\begin{aligned}
H(G) &= \frac{1}{2} \sum_{v \in V(G)} \widehat{D}_G(v) \leq \frac{1}{2} \left(\frac{5}{3}n^2 - n + \frac{2}{3} \sum_{i=1}^n (d(x_i) + d(y_i)) \right) \\
&\leq \frac{1}{2} \left(\frac{5}{3}n^2 - n + \frac{2}{3}(k^2 + (n-k)n + (n-k)^2 + kn) \right)
\end{aligned}$$

$$\begin{aligned} &\leq \frac{1}{2} \left(\frac{5}{3}n^2 - n + \frac{2}{3}((k + (n - k))^2 - 2k(n - k) + n^2) \right) \\ &\leq \frac{1}{2} \left(\frac{5}{3}n^2 - n + \frac{2}{3}(n^2 - 2 * 1 * 1 + n^2) \right) = \frac{9n^2 - 3n - 4}{6}. \end{aligned}$$

From $H(G) \geq \frac{9n^2 - 3n - 4}{6}$, $1 \leq k < n$, we have that $k = 1$, $n - k = 1$, $d(x_1) = 1$, $d(x_2) = 2$, $d(y_1) = 2$, and $d(y_2) = 1$. Thus $G = P_4$, which is not Hamiltonian.

This completes the proof of [Theorem 4](#). \square

Proof of Theorem 5. Let G be a graph satisfying the conditions in [Theorem 5](#). Notice that if $G = K_2 \vee ((2K_1) \cup K_{n-4})$, then $H(G) = \frac{n^2 - 3n + 7}{2}$. Suppose that G is not Hamiltonian and G is not $K_2 \vee ((2K_1) \cup K_{n-4})$. Then from [Lemma 4](#), we have that $e(G) \leq C(n - 2, 2) + 3$. Therefore

$$\begin{aligned} H(G) &\leq \frac{n(n-1)}{4} + \frac{1}{4} \sum_{v \in V(G)} d(v) \\ &\leq \frac{n(n-1)}{4} + \frac{1}{2}e(G) \leq \frac{n(n-1) + 2C(n-2, 2) + 6}{4} \leq \frac{n^2 - 3n + 6}{2}, \end{aligned}$$

which is a contradiction.

This completes the proof of [Theorem 5](#). \square

Proof of Theorem 6. Let G be a graph satisfying the conditions in [Theorem 6](#). Notice that if $G = K_3 \vee ((3K_1) \cup K_{n-6})$, then $H(G) = \frac{n^2 - 4n + 15}{2}$. Suppose that G is not Hamiltonian and G is not $K_3 \vee ((3K_1) \cup K_{n-6})$. Then from [Lemma 5](#), we have that $e(G) \leq C(n - 3, 2) + 8$. Therefore

$$\begin{aligned} H(G) &\leq \frac{n(n-1)}{4} + \frac{1}{4} \sum_{v \in V(G)} d(v) \\ &\leq \frac{n(n-1)}{4} + \frac{1}{2}e(G) \leq \frac{n(n-1) + 2C(n-3, 2) + 16}{4} \leq \frac{n^2 - 4n + 14}{2}, \end{aligned}$$

which is a contradiction.

This completes the proof of [Theorem 6](#). \square

Proof of Theorem 7. Let G be a graph satisfying the conditions in [Theorem 7](#). Suppose that G is not Hamiltonian. Then from [Lemma 6](#), we have that $e(G) \leq C(n, 2) - (k + 1)(n - k - 1)/2$. Therefore

$$\begin{aligned} H(G) &\leq \frac{n(n-1)}{4} + \frac{1}{4} \sum_{v \in V(G)} d(v) \\ &\leq \frac{n(n-1)}{4} + \frac{1}{2}e(G) \\ &\leq \frac{n(n-1) + 2C(n, 2) - (k + 1)(n - k - 1)}{4} \\ &\leq \frac{2n(n-1) - (k + 1)(n - k - 1)}{4}, \end{aligned}$$

which is a contradiction.

This completes the proof of [Theorem 7](#). \square

References

- [1] J.A. Bondy, U.S.R. Murty, *Graph Theory with Applications*, Macmillan, London and Elsevier, New York, 1976.
- [2] O. Ivanciuc, T.S. Balaban, A.T. Balaban, Reciprocal distance matrix, related local vertex invariants and topological indices, *J. Math. Chem.* 12 (1993) 309–318.
- [3] D. Plavšić, S. Nikolić, N. Trinajstić, Z. Mihalić, On the Harary index for the characterization of chemical graphs, *J. Math. Chem.* 12 (1993) 235–250.

- [4] H. Hua, M. Wang, On Harary index and traceable graphs, *MATCH Commun. Math. Comput. Chem.* 70 (2013) 297–300.
- [5] W. Byer, D. Smeltzer, Edge bounds in nonhamiltonian k -connected graphs, *Discrete Math.* 307 (2007) 1572–1579.
- [6] C. Berge, *Graphs and Hypergraphs*, American Elsevier Publishing Company, 1976.